

Koszul Duality and Cohomology: Generalized Syzygies for Commutative Koszul Algebras

Joint Work in Progress with Vassily Gorbounov (Aberdeen)
Zain Shaikh (Cologne) and Andrew Tonks (Londonmet)

Imma Gálvez Carrillo

UPC-EET, Terrassa

UAB Algebra and Combinatorics Seminar 2011

CRM

May 26, 2011

Towards generalized syzygies

Definition: Koszul homology

Let $\{a_1, \dots, a_n\}$ be a sequence of elements in a commutative \mathbb{C} -algebra A . Let W be an n -dimensional complex vector space with basis $\{\theta_1, \dots, \theta_n\}$. The **Koszul homology** of A with respect to the sequence $\{a_1, \dots, a_n\}$ is the homology of the complex

$$A \otimes \bigwedge W,$$

where A has homological degree zero, each θ_i has homological degree one, and the **Koszul differential** is given by the formula

$$d_K = \sum_{i=1}^n a_i \frac{\partial}{\partial \theta_i}.$$



Koszul Algebras (I)

Definition: Quadratic algebra

Let A be a positively graded connected algebra, locally finite-dimensional. A is called **quadratic** if it is determined by a vector space of generators $V = A_1$ and subspace of quadratic relations $I \subset A_1 \otimes A_1$

Definition: Koszul quadratic dual

The **Koszul dual** algebra $A^!$ associated with a quadratic algebra A is

$$A^! = T(V^*)/I^\perp,$$

where $I^\perp \subset V^* \otimes V^*$ is the annihilator of I . Clearly $A^{!!} = A$.

Koszul algebras (II)

Definition: Koszul algebra

A quadratic as above is a **Koszul algebra** iff

$$A^! \cong \text{Ext}_A^*(\mathbb{k}, \mathbb{k})$$

Lie superalgebras (I)

Definition: Lie superalgebra

A **Lie superalgebra** over \mathbb{C} is a $\mathbb{Z}/2$ -graded vector space (over \mathbb{C}) $L = L_{(0)} \oplus L_{(1)}$ with a map $[\cdot, \cdot] : L \otimes L \rightarrow L$ of $\mathbb{Z}/2$ -graded spaces, satisfying:

- 1 (anti-symmetry) $[x, y] = -(-1)^{|x||y|}[y, x]$ for all homogeneous $x, y \in L$,
- 2 (Jacobi identity) $(-1)^{|x||z|}[x, [y, z]] + (-1)^{|y||x|}[y, [z, x]] + (-1)^{|z||y|}[z, [x, y]] = 0$ for all homogeneous $x, y, z \in L$.

Lie superalgebras (II)

Definition: Lie superalgebra (continued)

Here $|x|$ is the parity of x : $|x| = i$ when $x \in L_{(i)}$ for $i = 0, 1$. An element x in $L_{(0)}$ or $L_{(1)}$ is termed even or odd respectively. We recover the familiar definition of a Lie algebra (over \mathbb{C}) in the case $L = L_{(0)}$.

Definition: graded Lie superalgebra

A **graded Lie superalgebra** is a Lie superalgebra L together with a grading compatible with the bracket and supergrading. That is, $L = \bigoplus_{m \geq 1} L_m$ such that $[L_i, L_j] \subset L_{i+j}$ and $L_{(i)} = \bigoplus_{m \geq 1} L_{2m-i}$ for $i = 0, 1$.

Koszul dual as universal envelope

Koszul dual as universal envelope

Assume that $A = T(V)/I$ is commutative, hence $\bigwedge^2 V \subset I$.

Therefore I^\perp is contained in $S^2(V^*)$, and so is generated by certain linear combinations of anti-commutators $[a_i^*, a_j^*] = a_i^* a_j^* + a_j^* a_i^*$.

As a consequence, the Koszul quadratic dual of A can be described as the universal envelope of a graded Lie superalgebra,

$$A^\natural = U(L), \quad L = \bigoplus_{m \geq 1} L_m = \mathbb{L}(V^*)/J, \quad (1)$$

where \mathbb{L} is the free Lie superalgebra functor, the space of (odd) generators V^* is concentrated in degree 1, and J is the Lie ideal with the same generators as I^\perp but viewed as linear combinations of supercommutators.

Some Lie ideals

The following Lie ideals are main characters of our work.

Definition

For $k \geq 2$, we define the graded Lie superalgebras

$$L_{\geq k} = \bigoplus_{m \geq k} L_m.$$

The algebra of syzygies (I)

Definition: the algebra of syzygies

Let A be a commutative \mathbb{C} -algebra which is a module over $S(V)$, and suppose we have a minimal free resolution of A ,

$$\cdots \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow A \rightarrow 0.$$

This is an exact sequence of graded free $S(V)$ -modules,

$$F_p = \bigoplus_{q \geq m_p} R_{pq} \otimes S(V),$$

where R_{pq} is the finite dimensional vector spaces of **p -th order syzygies of degree q** for A , and m_p is the minimum degree among the p -th order syzygies.

The algebra of syzygies (II)

Syzygies and the Koszul complex

- Since the chosen resolution is minimal, the differential vanishes on tensoring this complex with the trivial $S(V)$ -module \mathbb{C} .
- Hence

$$\mathrm{Tor}_p^{S(V)}(A, \mathbb{C}) = R_p := \bigoplus_{q \geq m_p} R_{pq}.$$

The algebra of syzygies (III)

- The functor $\text{Tor}^{S(V)}$ may also be calculated by resolving the other argument \mathbb{C} .
- The Koszul complex $K(S(V))$ of the symmetric algebra $S(V) = \mathbb{C}[a_1, \dots, a_n]$, is given by $(S(V) \otimes \wedge W, d_K)$, that is,

$$(\mathbb{C}[a_1, \dots, a_n] \otimes \wedge(\theta_1, \dots, \theta_n), d_K). \quad (2)$$

with the Koszul differential d_K of the sequence $\{a_1, \dots, a_n\}$.

The algebra of syzygies (IV)

Since this complex is a resolution of \mathbb{C} we can calculate the syzygies R_p of the quadratic algebra $A = T(V)/I$ by the homology of the complex

$$A \otimes_{S(V)} K(S(V)) = A \otimes_{S(V)} S(V) \otimes_{\mathbb{C}} \bigwedge W = A \otimes \bigwedge W. \quad (3)$$

This homology inherits a multiplication from $K(S(V))$ and becomes an associative algebra.

Lie ideals and the algebra of syzygies

- $H^*(L_{\geq 2}, \mathbb{C})$ gives indeed the algebra of syzygies of A .
- The interpretation of the algebras $L_{\geq k}$ for $k > 2$ was outlined by Berkovits in [4].
- We concentrate on the case $k = 3$.

Introduction: Berkovits' example

- In his work on string quantization and string/gauge theory duality [4], Berkovits uses crucially Koszul duality theory for commutative algebras.
- There, the coordinate algebra of the orthogonal Grassmannian $A = OGr(5, 10)$, related to the spinor representation of the group $SO(10, \mathbb{C})$ is considered.
- This algebra A is quadratic, and hence we know that its Koszul dual $A^!$ is $U(L)$ the universal enveloping algebra of a certain (graded) Lie superalgebra $L = \bigoplus_{i \geq 1} L_i$.

[4] Berkovits, N.; *Cohomology in the pure spinor formalism for the superstring*, J. High Energy Phys. 9 (2000).



- Berkovits' first observation, made in the context of string and gauge theories, was that the algebra of syzygies of A is isomorphic to the cohomology $H^*(L_{\geq 2}, \mathbb{C})$ of the Lie superalgebra $L_{\geq 2} = \bigoplus_{i \geq 2} L_i$.
- He proposed an extension of the Koszul complex of the coordinate algebra of $OGr(5, 10)$, relevant for his quantization procedure, and considered the question of calculating its homology.

- This extension is worthy of study for more general algebras.
- Berkovits' construction is an iteration of the Koszul homology of a sequence of elements of an algebra, first applied to the algebra A and then applied to its syzygies.
- As we recalled there is a notion of Koszul homology with respect to a sequence of elements $\{a_1, \dots, a_n\}$ of a commutative algebra A .
- And we recalled as well that if A is finitely generated and presented as $\mathbb{C}[a_1, \dots, a_n]/I$, the Koszul homology with respect to the sequence $\{a_i\}$ gives the algebra of syzygies of A .

- The generators Γ_j of the ideal I represent syzygies in the lowest degree.
- The Koszul homology of the algebra of syzygies with respect to the sequence Γ_j is the homology studied by Berkovits. This is what we call the algebra of generalized syzygies.
- It turns out that if A is Koszul, the generalized syzygies can also be described explicitly in terms of the Lie superalgebra L . This was shown by Movshev and Schwarz in [12] for the case of the coordinate algebra of the $OGr(5, 10)$.
- There was proved that in that case the homology of the Berkovits complex is isomorphic to the cohomology $H^*(L_{\geq 3}, \mathbb{C})$ of the Lie superalgebra $L_{\geq 3} = \bigoplus_{i \geq 3} L_i$.

[12] Movshev, M.; Schwarz, A.; *On maximally supersymmetric Yang-Mills theories*, Nuclear Physics B 681 (2004)



- Our aim is to extend these statements about $H^*(L_{\geq 3}, \mathbb{C})$ from the coordinate algebra of the orthogonal Grassmannian to an arbitrary finitely generated commutative Koszul algebra.
- Further examples are supplied by the orbits of the highest weight vector in an irreducible representation of a semisimple complex Lie group, as studied by Gorodentsev, Khoroshkin and Rudakov [9].
- We prove in our Main Theorem that the cohomology of the Lie algebra $L_{\geq 3}$ is isomorphic to the Berkovits homology of A .

[9] Gorodentsev, A.; Khoroshkin, A.; Rudakov A.; *On syzygies of highest weight orbits*, Amer. Math. Soc. Transl. 221 (2007).

- Since A is finitely generated, Hilbert's syzygy theorem says that the algebra of syzygies of A is finite dimensional.
- We expect the Lie superalgebra $L_{\geq k} \subset L_{\geq 3}$ to be a free Lie superalgebra for some k .
- Movshev and Schwarz, in [12], proved that $L_{\geq 3}$ is a free Lie superalgebra on an infinite set of generators in the case of the coordinate algebra of pure spinors.
- From the point of view of the gauge theory, freeness of $L_{\geq k}$ is important because it implies that the solution to the Euler-Lagrange equations for an abelian gauge group can be deformed to the solutions for a non-abelian gauge group.

- A commutative Koszul algebra A has a nice resolution in the category of algebras, given by the Chevalley complex of the Lie superalgebra L .
- Our main idea is to lift the Koszul differential which calculates the generalized syzygies to this resolution and subsequently calculate its homology.
- This requires a correction term to the naively lifted differential to guarantee that the lift squares to zero.
- This makes heavy use of the homological perturbation lemma à la Barnes and Lambe [3].

[3] Barnes, D.; Lambe, L.; *A fixed point approach to homological perturbation theory*, Proc. Amer. Math. Soc. 112 (1991), no. 3, 881–892.



Berkovits homology or generalized syzygies

Main Theorem

The algebra of generalized syzygies of a commutative Koszul algebra A is isomorphic to $H^*(L_{\geq 3}, \mathbb{C})$.

A result on DG algebras

Key Lemma

Let (C, d) be a commutative DG algebra over \mathbb{C} , nonnegatively graded and finitely generated in each degree. Let B be a contractible DG subalgebra of C , with quasi-isomorphism $\varepsilon : B \rightarrow \mathbb{C}$, and consider the DG ideal $\langle \overline{B} \rangle$ of C generated by the augmentation ideal $\overline{B} = \ker(\varepsilon)$. If $\langle \overline{B} \rangle$ is freely generated as a \overline{B} -module by a graded basis of homogeneous elements $Z = \bigcup_{i \geq 0} Z_i$, then C is quasi-isomorphic to $C/\langle \overline{B} \rangle$.

The Chevalley complex

Assume that A is a commutative Koszul algebra with $A^! = U(L)$ as above. We construct a resolution of A in the category of DG algebras from the Chevalley complex of L .

Definition: the Chevalley complex

The **Chevalley complex** of L is the cochain complex with

$$\mathrm{Ch}^i(L) = \left(\bigwedge^i L \right)^*$$

and the differential $d_C : \mathrm{Ch}^k(L) \rightarrow \mathrm{Ch}^{k+1}(L)$

$$(d_C \varphi)(x_0, \dots, x_k) = \sum_{i < j} (-1)^{j+\varepsilon(i,j)} \varphi(x_0, \dots, x_{i-1}, [x_i, x_j], x_{i+1}, \dots, \widehat{x}_j, \dots, x_k)$$



Some properties of the Chevalley complex

- $d_C : \text{Ch}^1(L) \rightarrow \text{Ch}^2(L)$ is the map that is dual to the bracket,

$$(d_C f)(x_0, x_1) = -f[x_0, x_1].$$

- The Chevalley complex is a cochain complex that calculates the cohomology of L with trivial coefficients, $H^*(L, \mathbb{C})$.
- This complex admits an algebra structure that descends to one on the cohomology of L so that the differential d_C is a derivation with respect to its product.

That is, given $\varphi \in (\wedge^i L)^*$ and $\psi \in (\wedge^j L)^*$,

$$\varphi \odot \psi \in (\wedge^{i+j} L)^* \text{ and } d_C(\varphi \odot \psi) = d_C\varphi \odot \psi \pm \varphi \odot d_C\psi.$$

The Chevalley complex as a chain complex

- As well as being a cochain complex whose cohomology is that of L , the Chevalley complex may also be considered a chain complex which defines a resolution of A .
- The chain complex is given by defining L_p^* to have homological grading $p - 1$, so that the homological and cohomological gradings together give the total degree in $\bigwedge L^*$.

The Chevalley complex as a chain complex (II)

We illustrate the Chevalley complex as a chain complex with homological grading as follows:

$$\mathrm{Ch}_3(L) \xrightarrow{d_C} \mathrm{Ch}_2(L) \xrightarrow{d_C} \mathrm{Ch}_1(L) \xrightarrow{d_C} \mathrm{Ch}_0(L) \longrightarrow 0$$

$$0 \longrightarrow L_1^* \longrightarrow 0$$

$$0 \longrightarrow L_2^* \longrightarrow \bigwedge^2 L_1^* \longrightarrow 0$$

$$0 \longrightarrow L_3^* \longrightarrow L_2^* \wedge L_1^* \longrightarrow \bigwedge^3 L_1^* \longrightarrow 0$$

The original cohomological grading is seen on the diagonals.

The Chevalley complex as resolution

In our situation of a graded Lie superalgebra L with $A^! = U(L)$ we observe that

$$\mathrm{Ch}_0(L) = \bigwedge L_1^* = S(V).$$

Proposition

For a commutative Koszul algebra A with $A^! = U(L)$, the chain complex given by the Chevalley complex of L with homological grading is a resolution of A .

The Chevalley complex as resolution (II)

Proof of proposition

The Chevalley complex of L with cohomological degree given by the number of exterior powers calculates

$$H^i(L, \mathbb{C})_j \cong \text{Ext}_{U(L)}^{ij}(\mathbb{C}, \mathbb{C}).$$

Since A is a commutative Koszul algebra, we have

$$A = \bigoplus_{i \geq 0} \text{Ext}_{A^!}^{ii}(\mathbb{C}, \mathbb{C})$$

and $\text{Ext}_{A^!}^{ij}(\mathbb{C}, \mathbb{C}) = 0$ for $i \neq j$

Chevalley complex as resolution (III)

Proof (continued)

But $A^! = U(L)$ and

$$H_i(\mathrm{Ch}(L))_j = H^{j-i}(L, \mathbb{C})_j,$$

where on the left hand side we are using our homological grading of $\mathrm{Ch}(L)$. Therefore,

$$H_i(\mathrm{Ch}(L)) = \begin{cases} A & \text{if } i = 0 \\ 0 & \text{if } i \neq 0 \end{cases}$$

and the Chevalley complex of L is a resolution of A .

Cohomology of $L_{\geq 2}$

Theorem

Let $A = T(V)/I$ be a commutative Koszul algebra, $R = \bigoplus_p R_p$ is its algebra of syzygies and $A^\dagger := T(V^*)/I^\perp = U(L)$ is its Koszul dual. Then $R_{pq} \cong H^{q-p}(L_{\geq 2}, \mathbb{C})_q$ as algebras.

Idea of proof

- Denote by X the tensor product $\text{Ch}(L) \otimes \bigwedge W$ with generators of W in degree one.
- Consider the differential $d_C + d_K$ on X , which one can check makes this into a complex.
- We put an algebra structure on the complex X by the following rule: if $\alpha, \beta \in \text{Ch}(L)$ and $\eta, \xi \in \bigwedge W$, then

$$(\alpha \otimes \eta) \cdot (\beta \otimes \xi) := (-1)^{|\eta||\beta|} \alpha \wedge \beta \otimes \eta \wedge \xi.$$

Idea of proof continued

- Since $\mathrm{Ch}_0(L) = S(V)$, this implies that $K(S(V)) = \mathrm{Ch}_0(L) \otimes \wedge W$ is a subalgebra in X .
- The restriction of the differential to this subalgebra is d_K , and hence is a resolution of \mathbb{C} .
- This satisfies the conditions of Key Lemma above and X is quasi-isomorphic to $X/\langle \overline{K(S(V))} \rangle$, where $\langle \overline{K(S(V))} \rangle$ is the DG ideal in X generated by the augmentation ideal of $K(S(V))$. Hence,
$$H_i(X) = H_i(X/\langle \overline{K(S(V))} \rangle) = H_i(\mathrm{Ch}(L_{\geq 2})),$$
 as $\mathrm{Ch}_0(L) = \mathrm{Ch}(L_1)$.

Idea of proof continued

- Now, consider the following filtration of X

$$\{0\} \subset F_0X \subset F_1X \subset \cdots \subset F_nX \subset \dots,$$

given by

$$F_pX_q := \sum_{j \leq p} \sum_{i+j=p+q} \text{Ch}_i(L) \otimes \bigwedge^j W.$$

- The differential on the E_0 -term of the spectral sequence associated to this filtration is d_C . Since $\bigwedge W$ is a vector space over \mathbb{C} , it is flat as a \mathbb{C} -module. Further, $\text{Ch}(L)$ is a resolution for A .
- We can conclude that the E_1 -term of the spectral sequence is contained in one line given by $A \otimes \bigwedge W$ with differential d_K .
- Hence, $R_i \cong H_i(X)$.

Idea of proof ended

- It is clear that both interpretations of the homology respect the multiplicative structure: $X/\langle\overline{K(S(V))}\rangle$ inherits its multiplicative structure from X and the spectral sequence will also respect it.
- The Chevalley complex $\text{Ch}(L)$ calculates the cohomology of Lie superalgebra L , where the cohomological grading of $\text{Ch}(L)$ is given by the number of exterior powers.
- Therefore, $R_{pq} = H_p(X)_q = H_p(\text{Ch}(L_{\geq 2}))_q = H^{q-p}(L_{\geq 2}, \mathbb{C})_q$.

Berkovits Complex

Let $A = \mathbb{C}[a_1, \dots, a_n]/I$ be commutative Koszul, with minimal set of generators $\{\Gamma_1, \dots, \Gamma_m\}$ of I representing lowest degree syzygies.

Lemma

If the quadratic relations for A are defined by the formulas

$$\Gamma_k = \sum_{i,j=1}^n \Gamma_{ij}^k a_i a_j,$$

for $k = 1, \dots, m$, then the representative for the homology class in the algebra of syzygies defined by the sequence $\{\Gamma_1, \dots, \Gamma_m\}$ is

$$\tilde{\Gamma}_k = \sum_{i,j=1}^n \Gamma_{ij}^k a_i \theta_j, \quad k = 1, \dots, m$$



Berkovits complex (II)

Definition

The **Berkovits complex** of a commutative Koszul algebra A is

$$A \otimes \bigwedge(\theta_1, \dots, \theta_n) \otimes \mathbb{C}[y_1, \dots, y_m]$$

equipped with the **Berkovits differential**

$$d_B = d_K + d_{Ber} = \sum_{i=1}^n a_i \frac{\partial}{\partial \theta_i} + \sum_{k=1}^m \sum_{i,j=1}^n \Gamma_{ij}^k a_i \theta_j \frac{\partial}{\partial y_k},$$

where the y_k have homological degree two.

(One can prove that $d_B^2 = 0$).

Lifting the differential

The setup

As before $L = \bigoplus L_i$ where $U(L) = A^!$. The vector space L_2 has the same dimension as $S^2(V^*)/I^\perp \cong I^*$ and the generators of I are linearly independent, so we have a basis $\{q_1, \dots, q_m\}$ of L_2^* such that

$$q_k = \sum_{i,j=1}^n \Gamma_{ij}^k \{a_i, a_j\},$$

and by construction,

$$d_C(q_k) = \sum_{i,j=1}^n \Gamma_{ij}^k a_i a_j$$

for $k = 1, \dots, m$.



Lifting the differential (II)

The idea

Define Y as:

$$\text{Ch}(L) \otimes \bigwedge(\theta_1, \dots, \theta_n) \otimes \mathbb{C}[y_1, \dots, y_m],$$

with an obvious graded algebra structure.

We can try to lift the Berkovits differential to Y as follows:

$d_C + d_K + d_{Ber}$ where

$$d_{Ber} = \sum_{i,j=1}^n \sum_{k=1}^m \Gamma_{ij}^k a_i \theta_j \frac{\partial}{\partial y_k}.$$

However, one checks that

$$(d_C + d_K + d_{Ber})^2 = \sum_{i,j=1}^n \sum_{k=1}^m \Gamma_{ij}^k a_i a_j \frac{\partial}{\partial y_k} \neq 0.$$

In order for the differential to square to zero, we define a correction to the differential as

$$d_S = - \sum_{k=1}^m q_k \frac{\partial}{\partial y_k}.$$

Proposition

$$(d_C + d_K + d_{Ber} + d_S)^2 = 0.$$

$$d_K d_S + d_S d_K = d_{Ber} d_S + d_S d_{Ber} = d_S^2 = 0,$$

and

$$d_C d_S + d_S d_C = - \sum_{i,j=1}^n \sum_{k=1}^m \Gamma_{ij}^k a_i a_j \frac{\partial}{\partial y_k} + \sum_{k=1}^m q_k \frac{\partial}{\partial y_k} d_C - \sum_{k=1}^m q_k \frac{\partial}{\partial y_k} d_C.$$

Therefore,

$$(d_C + d_K + d_{Ber} + d_S)^2 = (d_C + d_K + d_{Ber})^2 - \sum_{i,j=1}^n \sum_{k=1}^m \Gamma_{ij}^k a_i a_j \frac{\partial}{\partial y_k} = 0.$$

The Resolution of \mathbb{C} inside Y

Proposition

The subalgebra T of Y given by:

$$\text{Ch}(L_1, L_2) \otimes \bigwedge (\theta_1, \dots, \theta_n) \otimes \mathbb{C}[y_1, \dots, y_m]$$

is a resolution of \mathbb{C} .

This subcomplex is equipped with the differential

$$\sum_{i=1}^n a_i \frac{\partial}{\partial \theta_i} + \sum_{k=1}^m \left(\sum_{i,j=1}^n \Gamma_{ij}^k a_i \theta_j - q_k \right) \frac{\partial}{\partial y_k} + \sum_{k=1}^m \sum_{i,j=1}^n \Gamma_{ij}^k a_i a_j \frac{\partial}{\partial q_k}.$$

Idea of the proof

Step 1

The Koszul complex (P, d_K) of the sequence $\theta_1, \dots, \theta_n$ in $\text{Ch}(L_1)$ and the Koszul complex (Q, d_S) of y_1, \dots, y_m in $\text{Ch}(L_2)$ are contractible, as is the product $(P \otimes Q, d_K + d_S)$.

Step 2

We perturb the differential on $P \otimes Q$ to the differential on T by using the homological perturbation lemma of [3].

Step 3

The perturbed homotopy is well-defined, so T has a strong deformation retraction to $(\mathbb{C}, 0)$ concentrated in degree zero.

Step 1

Lemma

The complexes (P, d_K) and (Q, d_S) are contractible.
In fact, there are strong deformation retractions

$$\varphi_K \curvearrowright \mathbb{C}[a_1, \dots, a_n] \otimes \Lambda(\theta_1, \dots, \theta_n) \begin{array}{c} \xrightarrow{\epsilon} \\ \xleftarrow{\iota} \end{array} \mathbb{C}$$

$$\varphi_S \curvearrowright \Lambda(q_1, \dots, q_m) \otimes \mathbb{C}[y_1, \dots, y_m] \begin{array}{c} \xrightarrow{\epsilon} \\ \xleftarrow{\iota} \end{array} \mathbb{C}$$

where the right hand side is concentrated in degree zero, with trivial differential.

The same is true for the complex $(P \otimes Q, d_K + d_S)$.

Step 2

However, $\partial = d_K + d_S$ is not the differential we are interested in; we want to consider instead the complex with a perturbed differential,

$$(P \otimes Q, \partial + \partial').$$

Here ∂' is given by $d_C + d_{Ber}$,

$$\sum_{k=1}^m \sum_{i,j=1}^n \Gamma_{ij}^k a_i \theta_j \frac{\partial}{\partial y_k} + \Gamma_{ij}^k a_i a_j \frac{\partial}{\partial q_k}.$$

Then the complex is precisely T .

Step 2 continued

In order to show the complex with the new differential is also contractible we apply the homological perturbation lemma (see for example [3]). This says there is a unique perturbation of the deformation retraction

$$\varphi \curvearrowright (P \otimes Q, \partial) \begin{array}{c} \xrightarrow{\epsilon} \\ \xleftarrow{\iota} \end{array} \mathbb{C}$$

to a deformation retraction

$$\varphi' \curvearrowright (P \otimes Q, \partial + \partial') \begin{array}{c} \xrightarrow{\epsilon} \\ \xleftarrow{\iota} \end{array} \mathbb{C}$$

if and only if the original contracting homotopy φ and the perturbation of the differential ∂' satisfy a local nilpotence condition.

Step 2 ended

That is, for each element γ the expression

$$(\partial' \varphi)^N(\gamma)$$

is zero for sufficiently large $N = N(\gamma)$. In this case

$$\varphi \left(\sum_{n=0}^{\infty} (\partial' \varphi)^n \right)$$

is well-defined and will be the required contracting homotopy φ' .

Step 3

We proceed to prove the local nilpotency condition required.

Suppose $\alpha \otimes \beta \in P \otimes Q$ with $\beta \in \bigwedge^{\leq s}(q_1, \dots, q_m) \otimes \mathbb{C}[y_1, \dots, y_m]$.
If $s = 0$ then $\partial' \varphi(\alpha \otimes \beta) = 0$, and if $s > 0$ then

$$\partial' \varphi(\alpha \otimes \beta) \in P \otimes \bigwedge^{\leq (s-1)}(q_1, \dots, q_m) \otimes \mathbb{C}[y_1, \dots, y_m]$$

Hence $(\partial' \varphi)^{m+1} = 0$.

Generalized syzygies

Main Theorem

Let A be a commutative Koszul algebra and $A^! = U(L)$. Then,

$$H_{Ber}^*(A) \cong H^*(L_{\geq 3}, \mathbb{C}).$$

Idea of proof

- We have that T is a subalgebra in Y .
- We showed in a Proposition above that T is a resolution of \mathbb{C} . This satisfies the conditions of the Key Lemma and Y is quasi-isomorphic to $Y/\langle \overline{T} \rangle$, where $\langle \overline{T} \rangle$ is the DG ideal in Y generated by the augmentation ideal of T . Hence,
$$H_i(Y) = H_i(Y/\langle \overline{T} \rangle) = H_i(\text{Ch}(L_{\geq 3})).$$

Idea of proof, continued

- Now, consider the filtration of X

$$\{0\} \subset F_0X \subset F_1X \subset \cdots \subset F_nX \subset \cdots,$$

given by

$$F_p Y_q := \sum_{j \leq p} \sum_{i+j=p+q} \text{Ch}_i(L) \otimes \left(\bigwedge (\theta_1, \dots, \theta_n) \otimes \mathbb{C}[y_1, \dots, y_m] \right)_j.$$

- The differential on the E_0 -term of the spectral sequence associated to this filtration is d_C .
- Since, $\bigwedge (\theta_1, \dots, \theta_n) \otimes \mathbb{C}[y_1, \dots, y_m]$ is a vector space over \mathbb{C} , it is flat as a \mathbb{C} -module.





Idea of proof, ended







- As $\text{Ch}(L)$ is a resolution for A , we can conclude that the E_1 -term of the spectral sequence is contained in one line and is given by,




$$A \otimes \bigwedge (\theta_1, \dots, \theta_n) \otimes \mathbb{C}[y_1, \dots, y_m]$$

with precisely the Berkovits differential.

- Hence, the homology of this complex is also $H_{\text{Ber}}^*(A)$.
- The result follows by converting the grading to the cohomological grading of the Chevalley complex.

-  Aisaka, Y.; Kazama, Y.; *A New First Class Algebra, Homological Perturbation and Extension of Pure Spinor Formalism for Superstring*, UT-Komaba 02-15 hep-th/0212316 Dec, 2002.
-  Avramov, L.L.; *Free Lie subalgebras of the cohomology of local rings* Trans. Amer. Math. Soc. 270 (1982), no. 2, 589–608.
-  Barnes, D.; Lambe, L.; *A fixed point approach to homological perturbation theory*, Proc. Amer. Math. Soc. 112 (1991), no. 3, 881–892. See also Correction to: “A fixed point approach to homological perturbation theory”. Proc. Amer. Math. Soc. 129 (2001), no. 3, 941.
-  Berkovits, N.; *Cohomology in the pure spinor formalism for the superstring*, J. High Energy Phys. 9 (2000).

-  Chevalley, C.; Eilenberg, S.; *Cohomology Theory of Lie Groups and Lie Algebras* Trans. Amer. Math. Soc. 63, (1948). 85–124.
-  Eisenbud, D.; *Commutative Algebra with a View Toward Algebraic Geometry*, Grad. Text. in Math. Springer-Verlag, 1995. xvi+785 pp.
-  Gauss C.; *Disquisitiones generales de congruentis, Analysis residuorum. Caput octavum, Collected Works, Vol. 2*, Georg Olms Verlag, Hildersheim, New York, 1973, 212–242.
-  Gorbounov, V.; Schechtman, V.; *Divergent Series and Homological Algebra*, SIGMA 5 (2009).
-  Gorodentsev, A.; Khoroshkin, A.; Rudakov A.; *On syzygies of highest weight orbits*, Amer. Math. Soc. Transl. 221 (2007).
-  Hochschild, G.; *Relative homological algebra*, Trans. Amer. Math. Soc. 82 (1956), 246–269.

-  MacLane, S.; *Homology*, Reprint of the 1975 edition, Springer-Verlag, Berlin, 1995. x+422 pp.
-  Movshev, M.; Schwarz, A.; *On maximally supersymmetric Yang-Mills theories*, Nuclear Physics B 681 (2004)
-  Polishchuk, A.; Positselski, L.; *Quadratic algebras*, University Lecture Series, 37. AMS, (2005). xii+159 pp.