

# Non-persistence of resonant caustics in perturbed elliptic billiards

SÔNIA PINTO-DE-CARVALHO<sup>†</sup> and RAFAEL RAMÍREZ-ROS<sup>‡</sup>

<sup>†</sup> *Departamento de Matemática, ICEx, Universidade Federal de Minas Gerais,  
30.123–970, Belo Horizonte, Brazil*

*(e-mail: sonia@mat.ufmg.br)*

<sup>‡</sup> *Departament de Matemàtica Aplicada I, Universitat Politècnica de Catalunya,  
Diagonal 647, 08028 Barcelona, Spain*

*(e-mail: rafael.ramirez@upc.edu)*

*(Received 12 September 2011 and accepted in revised form 23 April 2012)*

*Abstract.* A caustic of a billiard table is a curve such that any billiard trajectory, once tangent to the curve, stays tangent after every reflection at the boundary. When the billiard table is an ellipse, any non-singular billiard trajectory has a caustic, which can be either a confocal ellipse or a confocal hyperbola. Resonant caustics—those whose tangent trajectories are closed polygons—are destroyed under generic perturbations of the billiard table. We prove that none of the resonant elliptical caustics persists under a large class of explicit perturbations of the original ellipse. This result follows from a standard Melnikov argument and the analysis of the complex singularities of certain elliptic functions.

## 1. Introduction and main result

Birkhoff [3] introduced the problem of *convex billiard tables* more than 80 years ago as a way to describe the motion of a free particle inside a closed convex smooth curve. The particle is reflected at the boundary according to the law that ‘the angle of incidence equals the angle of reflection’. Good modern starting points in the literature on the billiard problem are [14, 21]. Several open problems on billiard dynamics are listed in the survey [11].

A caustic is a curve such that any billiard trajectory, once tangent to the curve, stays tangent after every reflection. Caustics are the most distinctive geometric objects related to billiard tables, since they are a geometric manifestation of the regularity of their tangent trajectories. For example, integrable billiards have a continuum of caustics, whereas the non-existence of caustics inside a convex billiard table implies that there are some billiard trajectories whose past and future behaviours differ dramatically. See, for instance, [16]. Hence, the existence and persistence of caustics are two fundamental questions in billiards.

Most of the literature deals with convex caustics, since they are easier to understand and related to ordered trajectories. Two exceptions are [10, §3] and [13].

We summarize the classical existence results as follows. On the one hand, if the boundary curve is smooth enough and strictly convex, then there exists a collection of smooth convex caustics close to the boundary of the table whose union has positive area [7, 15]. On the other hand, Mather [16] proved that there are no smooth convex caustics inside a convex billiard table when its boundary curve has some flat point. Gutkin and Katok [10] gave a quantitative version of Mather's theorem.

The robustness of a smooth convex caustic is closely related to the arithmetic properties of its *rotation number*, which measures the number of turns around the caustic per impact. Caustics with Diophantine rotation numbers persist under small perturbations of the boundary curve. This follows from standard Kolmogorov–Arnold–Moser (KAM) arguments [7, 15]. On the contrary, *resonant caustics*—those whose tangent trajectories are closed polygons, so that their rotation numbers are rational—are fragile structures that generically break up. See, for instance, [19].

This raises two complementary questions: the first is to characterize the perturbations that preserve/destroy a given resonant caustic of a billiard table; the second is to determine all resonant caustics that are preserved/destroyed under a given perturbation of an integrable billiard table. These questions have been studied by several authors. Baryshnikov and Zharnitsky [2] proved that the perturbations preserving a given resonant caustic of a smooth convex billiard table form an infinite-dimensional Hilbert manifold. As a sample, we point out that this Hilbert manifold is given by the set of billiard tables with constant width when the rotation number of the unperturbed caustic is one half [13]. Concerning the second question, Ramírez-Ros [19] gave a sufficient condition for the break-up of the resonant circular caustics inside a circular billiard table, in terms of the Fourier coefficients of the perturbation, see Remark 3 below.

In this paper we tackle the second question when the billiard boundary is an ellipse. In that case, the billiard dynamics is integrable and any billiard trajectory has a caustic [21]. The caustics are the conics confocal to the original ellipse: confocal ellipses, confocal hyperbolas, and the foci. One can deduce from results of Poncelet [18] that if a billiard trajectory inside an ellipse is a closed polygon, then all the billiard trajectories sharing its caustic are also closed polygons. (We refer to [9] for a historical account of Poncelet's porisms.) Indeed, if a billiard trajectory tangent to one of the elliptical caustics is an  $(m, n)$ -gon—a closed polygon with  $n$  sides that makes  $m$  turns around its caustic—then all the billiard trajectories sharing its caustic are also  $(m, n)$ -gons, and their caustic is called  $(m, n)$ -resonant. (These two definitions are not restricted to billiards inside ellipses.) It is also known that there is a unique  $(m, n)$ -resonant elliptical caustic for any relatively prime integers  $m$  and  $n$  such that  $1 \leq m < n/2$ ; see §4. Our main result is that all these resonant elliptical caustics break up under a large class of explicit perturbations of the original ellipse; see Theorem 1.1.

The following notation is required to state the main result. Having fixed the ellipse

$$Q = \left\{ (x, y) \in \mathbb{R}^2 : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \right\}, \quad a > b > 0,$$

we consider its associated elliptic coordinates  $(\mu, \varphi)$  given by the relations

$$x = c \cosh \mu \cos \varphi, \quad y = c \sinh \mu \sin \varphi,$$

where  $c = \sqrt{a^2 - b^2}$  is the semifocal distance of  $Q$ . The equation of the ellipse  $Q$  in these elliptic coordinates is  $\mu \equiv \mu_0$ , where  $\cosh \mu_0 = a/c$  and  $\sinh \mu_0 = b/c$ . Hence, any smooth perturbation  $Q_\epsilon$  of the ellipse  $Q$  can be written in elliptic coordinates as

$$\mu = \mu_\epsilon(\varphi) = \mu_0 + \epsilon \mu_1(\varphi) + O(\epsilon^2), \tag{1}$$

for some  $2\pi$ -periodic smooth function  $\mu_\epsilon(\varphi)$ .

**THEOREM 1.1.** *Let  $\mu_1(\varphi)$  be a  $2\pi$ -periodic entire function. If  $\mu_1(\varphi)$  is not constant (respectively,  $\mu'_1(\varphi)$  is not  $\pi$ -antiperiodic), then none of the  $(m, n)$ -resonant elliptical caustics with odd  $n$  (respectively, even  $n$ ) persists under the perturbation (1).*

Our proof is based on the study of the persistence of the resonant rotational invariant circles (resonant RICs) of some twist maps by means of a first-order Melnikov method. Only convex caustics can be related to the RICs of those twist maps. Thus, there is no direct way to extend the same procedure to the non-convex caustic hyperbolas, but we believe that the same results hold for them.

*Remark 1.* If  $\mu_\epsilon(\varphi)$  is constant, then the perturbed curves  $Q_\epsilon$  are ellipses, so all caustics (resonant or not) are preserved. Hence, the hypothesis that  $\mu_1(\varphi)$  is non-constant is natural, since we are using a first-order method. Nevertheless, we can still state some results when this hypothesis fails. More precisely, let us assume that

$$\mu_\epsilon(\varphi) = \mu_0 + \epsilon \mu_1 + \dots + \epsilon^{i-1} \mu_{i-1} + \epsilon^i \mu_i(\varphi) + O(\epsilon^{i+1}),$$

for some  $\mu_0, \dots, \mu_{i-1} \in \mathbb{R}$  and some non-constant  $2\pi$ -periodic entire function  $\mu_i(\varphi)$ . Then:

- all the  $(m, n)$ -resonant elliptic caustics with odd  $n$  break up. This result is a corollary of Theorem 1.1. It suffices to consider  $\delta = \epsilon^i$  as the new perturbative parameter,  $Q_\epsilon^* = \{\mu \equiv \mu_0 + \dots + \epsilon^{i-1} \mu_{i-1}\}$  as the unperturbed ellipse, and to realize that  $Q_\epsilon$  is an  $O(\delta)$ -perturbation of  $Q_\epsilon^*$  whose first-order term in  $\delta$  satisfies the hypotheses of Theorem 1.1;
- if  $n$  is even, we believe that all  $(m, n)$ -resonant elliptical caustics also break up, even if  $\mu'_i(\varphi)$  is  $\pi$ -antiperiodic, but we should use a second-order Melnikov method in order to prove this. Unfortunately, the computations become too cumbersome.

*Remark 2.* If we write the perturbed ellipse  $Q_\epsilon$  in Cartesian coordinates as

$$x^2/a^2 + y^2/b^2 + \epsilon P_1(x, y) + O(\epsilon^2) = 1,$$

then

$$2(a^2 \sin^2 \varphi + b^2 \cos^2 \varphi) \mu_1(\varphi) + ab P_1(a \cos \varphi, b \sin \varphi) = 0.$$

In particular, the function  $\mu_1(\varphi)$  is  $\pi$ -antiperiodic when  $P_1(x, y)$  is odd.

*Remark 3.* The case of perturbed circular tables was studied using similar techniques in [19], but the final result was quite different. Let us recall it for comparison. Any billiard trajectory inside a circle of radius  $r_0$  has some concentric circle of radius  $\sqrt{r_0^2 - \lambda^2}$  as caustic, where  $0 < \lambda < r_0$  plays the role of a caustic parameter. If  $\lambda = r_0 \sin(m\pi/n)$ , then the circular caustic is  $(m, n)$ -resonant. Let us write the perturbed circle in polar coordinates  $(r, \theta)$  as

$$r = r_\epsilon(\theta) = r_0(1 + \epsilon r_1(\theta) + O(\epsilon^2)), \tag{2}$$

for some smooth function  $r_\epsilon : \mathbb{T} \rightarrow \mathbb{R}$ . Let  $\sum_{l \in \mathbb{Z}} \hat{r}_1^l e^{il\theta}$  be the Fourier expansion of  $r_1(\theta)$  and  $n \geq 2$ . If there exists some  $l \in n\mathbb{Z} \setminus \{0\}$  such that  $\hat{r}_1^l \neq 0$ , then the  $(m, n)$ -resonant circular caustics do not persist; see [19, Theorem 1]. In particular, it is not known if the  $(m, n)$ -resonant circular caustics with odd (respectively, even)  $n$  break up when  $r_1(\theta)$  is not constant (respectively,  $r_1'(\theta)$  is not  $\pi$ -antiperiodic).

In §2 we develop a general Melnikov theory to study the persistence of resonant RICs of twist maps. The general setup is adapted to billiard maps in §3. Finally, Theorem 1.1 is proved in §4 by analysing the complex singularities of certain elliptic functions, an idea borrowed from [6].

### 2. Break-up of resonant invariant curves in twist maps

This section is a generalization of [19, §2], since several hypotheses have been weakened. Namely, the unperturbed map can be non-integrable, the resonant invariant circle does not need to be horizontal, and the shift on the invariant circles can be non-constant. Nevertheless, the essential idea does not change. A similar theory is contained in [20]. For a general background on twist maps we refer to [12, §9.3] or [17].

Let  $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$ , and  $\pi_1 : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T}$  be the natural projection. Sometimes it is convenient to work in the universal cover  $\mathbb{R}$  of  $\mathbb{T}$ . We will use the coordinates  $(x, y)$  for both  $\mathbb{T} \times \mathbb{R}$  and  $\mathbb{R}^2$ . The lines of the form  $x = \text{constant}$  and  $y = \text{constant}$  will be called vertical and horizontal, respectively. A tilde will always denote the lift of a function or set to the universal cover. If  $g$  is a real-valued function,  $\partial_i g$  denotes the derivative with respect to the  $i$ th variable. We will assume that all the objects considered are smooth. Here, ‘smooth’ means  $C^\infty$ . In particular, all the dependences on the perturbative parameter  $\epsilon$  are assumed to be smooth.

We will consider certain diffeomorphisms defined on an open cylinder of the form  $Z = \mathbb{T} \times Y$ , for some open bounded interval  $Y = (y_-, y_+) \subset \mathbb{R}$ . Then  $\tilde{Z} = \mathbb{R} \times Y$  is an open strip of the plane. A diffeomorphism  $f : Z \rightarrow Z$  is called an *area-preserving twist map* when it preserves area and orientation and satisfies the *twist condition*

$$\partial_2 \tilde{\pi}_1 \tilde{f}(x, y) \neq 0 \quad \text{for all } (x, y) \in \tilde{Z}.$$

If the twist is positive (respectively, negative), then the first iterate of any vertical line tilts to the right (respectively, left). We also assume, although it is not essential, that  $f$  satisfies some *rigid boundary conditions*. To be more precise, we suppose that the twist map  $f$  can be extended continuously to the closed cylinder  $\mathbb{T} \times [y_-, y_+]$  as a rigid rotation on the boundaries. That is, there exist some *boundary frequencies*  $\omega_\pm \in \mathbb{R}$ ,  $\omega_- < \omega_+$ , such that  $\tilde{f}(x, y_\pm) = (x + \omega_\pm, y_\pm)$ .

Let

$$D = \{(x, x') \in \mathbb{R}^2 : \omega_- < x' - x < \omega_+\}.$$

Then there exists a function  $h : D \rightarrow \mathbb{R}$  such that  $\tilde{f}(x, y) = (x', y')$  if and only if

$$y = -\partial_1 h(x, x'), \quad y' = \partial_2 h(x, x'). \tag{3}$$

The function  $h$  is called the *generating function* of  $f$ . Furthermore, if  $(x'', y'') = \tilde{f}(x', y')$ , then

$$\partial_2 h(x, x') + \partial_1 h(x', x'') = 0. \tag{4}$$

We study the dynamics of  $f$ , but it is often more convenient to work with the lift  $\tilde{f}$ , so we will pass between the two without comment and, in what follows, the lift  $\tilde{f}$  remains fixed.

A closed curve  $\Upsilon \subset Z$  is said to be a *rotational invariant circle* (RIC) of  $f$  when it is homotopically non-trivial and  $f(\Upsilon) = \Upsilon$ . Birkhoff proved that all RICs are graphs of Lipschitz functions. See, for instance, [17, §IV.C]. Let  $v : \mathbb{T} \rightarrow Y$  be the Lipschitz function such that  $\Upsilon = \text{graph } v := \{(x, v(x)) : x \in \mathbb{T}\}$ . If  $v$  is smooth, we say that  $\Upsilon$  is a *smooth RIC*.

Twist maps do not form a closed set under composition. For instance, the square of a twist map is not necessarily a twist map, and indeed typically it is not. Nevertheless, any power of a twist map is *locally twist* on its smooth RICs.

LEMMA 2.1. *If  $\Upsilon = \text{graph } v$  is a smooth RIC of an area-preserving twist map  $f : Z \rightarrow Z$ , then*

$$\partial_2 \tilde{\pi}_1 \tilde{f}^n(x, \tilde{v}(x)) \neq 0 \quad \text{for all } x \in \mathbb{R}, \text{ for all } n \geq 1.$$

*Proof.* Given any point

$$p = (x, \tilde{v}(x)) \in \tilde{\Upsilon},$$

let

$$p_j = (x_j, \tilde{v}(x_j)) = \tilde{f}^j(p), \quad t_j = (1, \tilde{v}'(x_j)), \quad \text{and} \quad v_j = (0, 1).$$

We identify the tangent planes  $T_p \tilde{Z}$  with the Euclidean plane  $\mathbb{R}^2$ . Thus, the vector  $t_j$  is tangent to  $\tilde{\Upsilon}$  at the point  $p_j$  and  $v_j$  is a vertical vector at  $p_j$ . The linear map  $d\tilde{f}^n(p) : T_p \tilde{Z} \rightarrow T_{p_n} \tilde{Z}$  is the composition of the linear maps  $d\tilde{f}(p_j) : T_{p_j} \tilde{Z} \rightarrow T_{p_{j+1}} \tilde{Z}$  for  $j = 0, \dots, n - 1$ . Let  $a_j, b_j, c_j, d_j, \alpha_n, \beta_n, \gamma_n, \delta_n \in \mathbb{R}$  be the coefficients such that

$$\begin{aligned} d\tilde{f}(p_j) : t_j &\mapsto a_j t_{j+1} + c_j v_{j+1}, & v_j &\mapsto b_j t_{j+1} + d_j v_{j+1}, \\ d\tilde{f}^n(p) : t_0 &\mapsto \alpha_n t_n + \gamma_n v_n, & v_0 &\mapsto \beta_n t_n + \delta_n v_n. \end{aligned}$$

We note that

$$b_j = \partial_2 \tilde{\pi}_1 \tilde{f}(p_j) \quad \text{and} \quad \beta_n = \partial_2 \tilde{\pi}_1 \tilde{f}^n(p).$$

Let us suppose that the twist is positive, so  $b_j > 0$ . We wish to prove that  $\beta_n > 0$  for any integer  $n \geq 1$ . The case of negative twist is completely analogous.

We deduce that  $c_j = 0$  from the invariance of  $\tilde{\Upsilon}$ . Hence,  $\beta_n = \sum_{j=0}^{n-1} D_0^{j-1} b_j A_{j+1}^{n-1}$ , where  $D_i^j = \prod_{k=i}^j d_k$  and  $A_i^j = \prod_{k=i}^j a_k$ . Furthermore, we note that  $d_j > 0$  because the two components of  $C \setminus \Upsilon$  are invariant. Finally, we get that  $a_j > 0$  from the preservation of orientation. □

Roughly speaking, an RIC is said to be *resonant* when all its points are periodic, but we need to be more precise. Let  $(x, y) \in Z$  be a periodic point of the twist map  $f$ , and let  $n$  be its smallest period. Then, there exists an integer  $m$  such that its lift satisfies  $\tilde{f}^n(x, y) = (x + 2\pi m, y)$ . Obviously,  $\omega_- < 2\pi m/n < \omega_+$ . Such a periodic point is said to be of *type*  $(m, n)$ . An RIC is said to be  $(m, n)$ -*resonant* when all its points are periodic of type  $(m, n)$ .

Let  $f$  be an area-preserving twist map with an  $(m, n)$ -resonant smooth RIC  $\Upsilon = \text{graph } \nu$ . Considering area-preserving twist perturbations of the form  $f_\epsilon = f + O(\epsilon)$ , we prove in the following lemma that there exist two graphs  $\Upsilon_\epsilon = \text{graph } \nu_\epsilon$  and  $\Upsilon_\epsilon^* = \text{graph } \nu_\epsilon^*$   $O(\epsilon)$ -close to  $\Upsilon$  and such that  $f_\epsilon^n$  projects the first graph onto the second along the vertical direction.

LEMMA 2.2. *There exist two smooth functions  $\nu_\epsilon, \nu_\epsilon^* : \mathbb{T} \rightarrow Y$  defined for  $\epsilon \in (-\epsilon_0, \epsilon_0)$ ,  $\epsilon_0 > 0$ , such that:*

- (1)  $\nu_\epsilon(x) = \nu(x) + O(\epsilon)$  and  $\nu_\epsilon^*(x) = \nu(x) + O(\epsilon)$ , uniformly in  $x \in \mathbb{T}$ ; and
- (2)  $f_\epsilon^n(x, \nu_\epsilon(x)) = (x, \nu_\epsilon^*(x))$ , for all  $x \in \mathbb{T}$ .

*Proof.* We work with the lift of the maps. Having fixed an angle  $x \in \mathbb{R}$ , let  $y_0 = \tilde{\nu}(x)$  and

$$\tilde{G}(y, \epsilon) := \tilde{\pi}_1 \tilde{f}_\epsilon^n(x, y) - x - 2\pi m.$$

This function  $\tilde{G}(y, \epsilon)$  satisfies the hypotheses of the implicit function theorem at the point  $(y, \epsilon) = (y_0, 0)$ , since  $\tilde{G}(y_0, 0) = 0$  and  $\partial_1 \tilde{G}(y_0, 0) = \partial_2 \tilde{\pi}_1 \tilde{f}_\epsilon^n(x, \tilde{\nu}(x)) \neq 0$ ; see Lemma 2.1. Consequently, there exist  $\epsilon_0, \eta > 0$  such that the equation  $\tilde{G}(y, \epsilon) = 0$  has exactly one solution  $y_\epsilon = y_0 + O(\epsilon)$  in the interval  $(y_0 - \eta, y_0 + \eta)$  for all  $\epsilon \in (-\epsilon_0, \epsilon_0)$ . We recall that  $\tilde{G}(y, \epsilon)$  had  $x \in \mathbb{R}$  as an extra parameter, but it appeared in a  $2\pi$ -periodic smooth way. Hence,  $\epsilon_0$  and  $\eta$  can be taken independent of  $x$ , the estimate  $|y_\epsilon - y_0| = O(\epsilon)$  is uniform in  $x$ , and  $y_\epsilon$  depends in a  $2\pi$ -periodic smooth way on  $x$ . Finally, set  $\tilde{\nu}_\epsilon(x) = y_\epsilon$  and then  $\tilde{\nu}_\epsilon^*(x)$  is determined by means of the relation  $\tilde{f}_\epsilon^n(x, \tilde{\nu}_\epsilon(x)) = (x + 2\pi m, \tilde{\nu}_\epsilon^*(x))$ . The functions  $\tilde{\nu}_\epsilon, \tilde{\nu}_\epsilon^* : \mathbb{R} \rightarrow Y$  are  $2\pi$ -periodic and smooth, so they can be projected to two smooth functions  $\nu_\epsilon, \nu_\epsilon^* : \mathbb{T} \rightarrow Y$  that satisfy the two claimed properties by construction. □

We say that an  $(m, n)$ -resonant smooth RIC  $\Upsilon$  of a twist map  $f$  *persists* under an area-preserving twist perturbation  $f_\epsilon = f + O(\epsilon)$  whenever the perturbed map has an  $(m, n)$ -resonant RIC  $\Upsilon_\epsilon$  for any small enough  $\epsilon$  such that  $\Upsilon_\epsilon = \Upsilon + O(\epsilon)$ . The corollary below follows immediately from this definition.

COROLLARY 2.1. *The resonant RIC  $\Upsilon$  persists under the perturbation  $f_\epsilon$  if and only if  $\Upsilon_\epsilon = \Upsilon_\epsilon^*$ .*

Therefore, it is quite useful to quantify the separation between the graphs  $\Upsilon_\epsilon$  and  $\Upsilon_\epsilon^*$ .

LEMMA 2.3.  $\nu_\epsilon^*(x) - \nu_\epsilon(x) = L'_\epsilon(x)$ , where  $L_\epsilon : \mathbb{T} \rightarrow \mathbb{R}$  is a function whose lift is

$$\tilde{L}_\epsilon(x) = \sum_{j=0}^{n-1} h_\epsilon(\bar{x}_j(x; \epsilon), \bar{x}_{j+1}(x; \epsilon)), \quad \bar{x}_j(x; \epsilon) = \tilde{\pi}_1 \tilde{f}_\epsilon^j(x, \tilde{\nu}_\epsilon(x)), \tag{5}$$

and  $h_\epsilon$  is the generating function of  $f_\epsilon$ .

*Proof.* As long as no confusion results, we will omit the dependence on  $x$  and  $\epsilon$ . We write  $(\bar{x}_j, \bar{y}_j) = \tilde{f}^j(x, \tilde{v}(x))$  and  $\bar{w}_j = \partial \bar{x}_j / \partial x$  for  $j = 0, \dots, n$ . Then  $\bar{x}_0 = x$  and  $\bar{x}_n = x + 2\pi m$ , so  $\bar{w}_0 = \bar{w}_n = 1$ . Furthermore,  $\bar{y}_0 = \tilde{v}(x)$  and  $\bar{y}_n = \tilde{v}^*(x)$ . From the implicit equations (3), we get that  $\partial_1 h(\bar{x}_0, \bar{x}_1) = -\bar{y}_0$ ,  $\partial_2 h(\bar{x}_{n-1}, \bar{x}_n) = \bar{y}_n$ , and  $\partial_2 h(\bar{x}_{j-1}, \bar{x}_j) + \partial_1 h(\bar{x}_j, \bar{x}_{j+1}) = 0$  for  $j = 1, \dots, n - 1$ . Therefore,

$$\begin{aligned} \tilde{L}'(x) &= \partial_1 h(\bar{x}_0, \bar{x}_1)\bar{w}_0 + \sum_{j=1}^{n-1} (\partial_2 h(\bar{x}_{j-1}, \bar{x}_j) + \partial_1 h(\bar{x}_j, \bar{x}_{j+1}))\bar{w}_j \\ &\quad + \partial_2 h(\bar{x}_{n-1}, \bar{x}_n)\bar{w}_n = \tilde{v}^*(x) - \tilde{v}(x). \end{aligned}$$

It is simple to check that  $\tilde{L} : \mathbb{R} \rightarrow \mathbb{R}$  is  $2\pi$ -periodic, so it can be projected to a function  $L : \mathbb{T} \rightarrow \mathbb{R}$ . □

**COROLLARY 2.2.** *The resonant RIC  $\Upsilon$  persists under the perturbation  $f_\epsilon$  if and only if  $L'_\epsilon(x) \equiv 0$ .*

We shall say that  $L_\epsilon : \mathbb{T} \rightarrow \mathbb{R}$  is the *subharmonic potential* of the resonant RIC  $\Upsilon$  under the twist perturbation  $f_\epsilon$ . It is quite natural to extract information from the low-order terms of its expansion  $L_\epsilon(x) = L_0(x) + \epsilon L_1(x) + O(\epsilon^2)$ . This is the main idea behind any Melnikov approach to a perturbative problem. The zero-order term  $L_0(x)$  is constant (and so of no use), since  $L'_0(x) = \nu_0^*(x) - \nu_0(x) = \nu(x) - \nu(x) \equiv 0$ . We shall say that the first-order term  $L_1(x)$  is the *subharmonic Melnikov potential* of the resonant RIC  $\Upsilon$  under the twist perturbation  $f_\epsilon$ . The proposition below provides a closed formula for its computation.

**PROPOSITION 2.1.** *If  $h_\epsilon = h + \epsilon h_1 + O(\epsilon^2)$ , then the lift of  $L_1(x)$  is*

$$\tilde{L}_1(x) = \sum_{j=0}^{n-1} h_1(x_j, x_{j+1}), \quad x_j = \tilde{\pi}_1 \tilde{f}^j(x, \tilde{v}(x)).$$

*Proof.* Given any  $x \in \mathbb{R}$ , we set  $x_j = x_j(x) := \bar{x}_j(x; 0)$  and  $z_j = z_j(x) := \partial_2 \bar{x}_j(x; 0)$  for  $j = 0, \dots, n$ . Then the  $O(\epsilon)$ -term of (5) is

$$\begin{aligned} \tilde{L}_1(x) &= \partial_1 h(x_0, x_1)z_0 + \sum_{j=1}^{n-1} (\partial_1 h(x_j, x_{j+1}) + \partial_2 h(x_{j-1}, x_j))z_j \\ &\quad + \partial_2 h(x_{n-1}, x_n)z_n + \sum_{j=0}^{n-1} h_1(x_j, x_{j+1}). \end{aligned}$$

Using the implicit equations (3) for the unperturbed twist map, the first summation vanishes. The terms  $\partial_1 h(x_0, x_1)z_0$  and  $\partial_2 h(x_{n-1}, x_n)z_n$  also vanish, since

$$\bar{x}_0(x; \epsilon) = x \quad \text{and} \quad \bar{x}_n(x; \epsilon) = x + 2\pi m \quad \text{for all } \epsilon \in (-\epsilon_0, \epsilon_0).$$

Furthermore,

$$x_j = x_j(x) = \bar{x}_j(x; 0) = \tilde{\pi}_1 \tilde{f}^j(x, \nu(x)). \quad \square$$




FIGURE 1. The billiard map  $f(\varphi, \vartheta) = (\varphi', \vartheta')$ .

The following corollary displays the most important property of the subharmonic Melnikov potential in relation to the goals of this paper.

**COROLLARY 2.3.** *If  $L_1(x)$  is not constant, then the resonant RIC  $\Upsilon$  does not persist under the perturbation  $f_\epsilon$ .*

*Proof.* This follows directly from Corollary 2.2 and the estimate  $L_\epsilon = \text{constant} + \epsilon L_1 + O(\epsilon^2)$ . □

### 3. Break-up of resonant caustics in perturbed billiard tables

Let  $Q$  be a closed strictly convex smooth curve in the plane. Let  $\gamma : \mathbb{T} \rightarrow Q$  be a counterclockwise parametrization. Let  $Z = \mathbb{T} \times (0, \pi)$  be an open cylinder. We can model the billiard dynamics inside  $Q$  by means of a map  $f : Z \rightarrow Z$ ,  $f(\varphi, \vartheta) = (\varphi', \vartheta')$ , defined as follows. If the particle hits  $Q$  at a point  $\gamma(\varphi)$  under an angle of incidence  $\vartheta \in (0, \pi)$  with the tangent vector at  $\gamma(\varphi)$ , then, as the motion is free inside  $Q$ , the next impact point is  $\gamma(\varphi')$ , the intersection point with the boundary, and the next angle of incidence is  $\vartheta' \in (0, \pi)$ , as in Figure 1. A straightforward computation shows that  $f(\varphi, \vartheta) = (\varphi', \vartheta')$  if and only if

$$|\gamma'(\varphi)| \cos \vartheta = -\partial_1 h(\varphi, \varphi'), \quad |\gamma'(\varphi')| \cos \vartheta' = \partial_2 h(\varphi, \varphi'), \tag{6}$$

where  $h : \mathbb{T}^2 \setminus \{\varphi' \neq \varphi\} \rightarrow \mathbb{R}$  is given by  $h(\varphi, \varphi') = |\gamma(\varphi) - \gamma(\varphi')|$ . Furthermore, the twist condition holds:  $\partial\varphi'/\partial\vartheta = h(\varphi, \varphi')/|\gamma'(\varphi')| \sin \vartheta' > 0$ . Finally, it is geometrically clear that  $f$  satisfies the rigid boundary conditions with  $\omega_- = 0$  and  $\omega_+ = 2\pi$ .

A remark is in order. Equations (6) differ slightly from equations (3), but identity (4) still holds and so the theory developed in the previous section still applies.

Obviously, one could write the map in the canonical coordinates—arc length parameter for the boundary and  $\cos \vartheta$  as its conjugate—in order to have  $h$  as a generating function, but this is not a wise choice when dealing with ellipses.

Let us assume that there exists a closed convex smooth caustic  $C$  contained in the region enclosed by  $Q$ . Then the billiard map  $f : Z \rightarrow Z$  has two smooth RICs  $\Upsilon^\pm = \text{graph } \vartheta^\pm \subset Z$ . The functions  $\vartheta^\pm : \mathbb{T} \rightarrow (0, \pi)$  are easy to understand:  $\vartheta^+(\varphi)$  and  $\vartheta^-(\varphi)$






FIGURE 2. Left: A (1, 4)-resonant convex smooth caustic  $C$ . Right: Its two smooth RICs  $\Upsilon^- = \text{graph } \vartheta^-$  and  $\Upsilon^+ = \text{graph } \vartheta^+$  in the phase space  $Z = \mathbb{T} \times (0, \pi)$ .

are the angles determined by the two lines tangent to the caustic  $C$  from the point  $\gamma(\varphi) \in Q$ ; see Figure 2. In particular,  $\vartheta^-(\varphi) + \vartheta^+(\varphi) = \pi$ . To fix ideas, we will assume that  $\Upsilon^-$  and  $\Upsilon^+$  correspond to the billiard motion around  $C$  in the counterclockwise and clockwise senses, respectively. Hence,  $0 < \vartheta^-(\varphi) < \pi/2 < \vartheta^+(\varphi) < \pi$ . There is an explicit formula relating the parametrization of the billiard curve  $Q$ , the parametrization of the caustic  $C$ , and the functions  $\vartheta^\pm$ . See, for instance, [7, 13].

Let  $Q$  be a closed strictly convex smooth billiard boundary with an  $(m, n)$ -resonant convex caustic  $C$ , so that its RIC  $\Upsilon^-$  is  $(m, n)$ -resonant and its RIC  $\Upsilon^+$  is  $(n - m, n)$ -resonant. We say that  $C$  persists under a perturbation  $Q_\epsilon = Q + O(\epsilon)$  whenever the perturbed billiard curve has an  $(m, n)$ -resonant caustic  $C_\epsilon$  for any small enough  $\epsilon$  such that  $C_\epsilon = C + O(\epsilon)$ .

Let  $f_\epsilon$  be the billiard map inside  $Q_\epsilon$  and  $L_1^-(\varphi)$  and  $L_1^+(\varphi)$  be the subharmonic Melnikov potentials of the resonant RICs  $\Upsilon^-$  and  $\Upsilon^+$  under the area-preserving twist perturbation  $f_\epsilon$ . Both potentials coincide, due to the time reversibility of the billiard dynamics. Therefore, we can omit the  $\pm$  signs. In this context, we will say that  $L_1(\varphi)$  is the subharmonic Melnikov potential of the resonant caustic  $C$  for the perturbation  $Q_\epsilon$ .

**COROLLARY 3.1.** *If  $L_1(\varphi)$  is not constant, then the resonant caustic  $C$  does not persist under the perturbation  $Q_\epsilon$ .*

4. Break-up of resonant caustics in perturbed elliptic billiard tables

From now on, we will assume that the unperturbed billiard boundary is the ellipse

$$Q = \left\{ q = (x, y) \in \mathbb{R}^2 : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \right\}, \quad a > b > 0.$$

It is known that the convex caustics of the billiard inside  $Q$  are the confocal ellipses

$$C_\lambda = \left\{ q = (x, y) \in \mathbb{R}^2 : \frac{x^2}{a^2 - \lambda^2} + \frac{y^2}{b^2 - \lambda^2} = 1 \right\}, \quad 0 < \lambda < b.$$

Let  $\rho(\lambda)$  be the rotation number of the elliptical caustic  $C_\lambda$ . In [4, 8] it is proved that  $\rho : (0, b) \rightarrow \mathbb{R}$  is an analytic increasing function such that  $\rho(0) = 0$  and  $\rho(b) = 1/2$ .

The second reference is based on the study of Fomenko graphs, whereas the first one uses the theory of ordered circle homeomorphisms. Thus, there is a unique  $(m, n)$ -resonant elliptical caustic for any relatively prime integers  $m$  and  $n$  such that  $1 \leq m < n/2$ . We shall see that the caustic parameter  $\lambda \in (0, b)$  of the  $(m, n)$ -resonant caustic is implicitly determined by means of an equation containing a couple of elliptic integrals; see equation (10).

The following lemma on elliptic billiards is useful to simplify the expression for the subharmonic Melnikov potential later on.

LEMMA 4.1. *Let  $(q_j)_{j \in \mathbb{Z}}$  be any billiard trajectory inside the ellipse  $Q$  with caustic  $C_\lambda$ . Let  $p_j = (q_{j+1} - q_j)/|q_{j+1} - q_j|$  be the unit inward velocities of the trajectory. Then*

$$ab \langle p_{j-1} - p_j, D^{-2}q_j \rangle = 2\lambda \quad \text{for all } j \in \mathbb{Z},$$

where  $D = \text{diag}(a, b)$  is the diagonal matrix such that  $Q = \{q \in \mathbb{R}^2 : \langle q, D^{-2}q \rangle = 1\}$ .

*Proof.* We shall prove that, given any point  $q = (x, y) \in Q$  and any unit inward vector  $p = (u, v) \in \mathbb{S}^1$ , the line  $\ell = \{q + \tau p : \tau \in \mathbb{R}\}$  is tangent to the conic  $C_\lambda$  if and only if

$$\lambda = -(bxu/a + ayv/b) = -ab \langle p, D^{-2}q \rangle.$$

To begin with, we note that the line  $\ell$  is tangent to the conic  $C_\lambda$  if and only if the equation of second order in the variable  $\tau$  given by

$$(x + \tau u)^2 / (a^2 - \lambda^2) + (y + \tau v)^2 / (b^2 - \lambda^2) - 1 = 0$$

has zero discriminant, which is equivalent to the equation

$$\left( \frac{xu}{a^2 - \lambda^2} + \frac{yv}{b^2 - \lambda^2} \right)^2 = \left( \frac{u^2}{a^2 - \lambda^2} + \frac{v^2}{b^2 - \lambda^2} \right) \left( \frac{x^2}{a^2 - \lambda^2} + \frac{y^2}{b^2 - \lambda^2} - 1 \right).$$

After some simplifications, we can rewrite this equation as

$$(xv - yu)^2 = (b^2 - \lambda^2)u^2 + (a^2 - \lambda^2)v^2 = a^2v^2 + b^2u^2 - \lambda^2,$$

since  $u^2 + v^2 = 1$ . Next, using the fact that  $x^2/a^2 + y^2/b^2 = 1$ , we obtain that

$$\lambda^2 = (a^2v^2 + b^2u^2)(x^2/a^2 + y^2/b^2) - (xv - yu)^2 = (bxu/a + ayv/b)^2.$$

Thus, we have two possibilities:  $\lambda = ab \langle p, D^{-2}q \rangle$  or  $\lambda = -ab \langle p, D^{-2}q \rangle$ . The first is discarded because  $\lambda > 0$  and  $\langle p, D^{-2}q \rangle < 0$ . The second inequality follows from the fact that the vector  $p$  points inward  $Q$  at  $q$ , whereas  $D^{-2}q$  is an outward normal vector to  $Q$  at  $q$ .

Finally, we note that  $-p_{j-1} = (q_{j-1} - q_j)/|q_{j-1} - q_j|$  and  $p_j = (q_{j+1} - q_j)/|q_{j+1} - q_j|$  are the two unit vectors that point inward  $Q$  at the impact point  $q_j$  and give the two tangent directions to the caustic  $C_\lambda$ . Therefore,  $\lambda = ab \langle p_{j-1}, D^{-2}q_j \rangle = -ab \langle p_j, D^{-2}q_j \rangle$ .  $\square$

PROPOSITION 4.1. *Let  $C_\lambda$  be the  $(m, n)$ -resonant elliptical caustic confocal to the ellipse  $Q$ . Given any angle  $\varphi \in \mathbb{T}$ , let  $q_j = (a \cos \varphi_j, b \sin \varphi_j)$  be the vertices of the*

( $m, n$ )-gon inscribed in  $Q$  and circumscribed around  $C_\lambda$  such that  $q_0 = (a \cos \varphi, b \sin \varphi)$ . Then the subharmonic Melnikov potential of the caustic  $C_\lambda$  for the perturbed ellipse (1) is

$$L_1(\varphi) = 2\lambda \sum_{j=0}^{n-1} \mu_1(\varphi_j). \tag{7}$$

*Proof.* The parametrization of the perturbed ellipse (1) is given by

$$\gamma_\epsilon(\varphi) = (c \cosh \mu_\epsilon(\varphi) \cos \varphi, c \sinh \mu_\epsilon(\varphi) \sin \varphi) = \gamma_0(\varphi) + \epsilon \gamma_1(\varphi) + O(\epsilon^2),$$

where  $\gamma_0(\varphi) = (a \cos \varphi, b \sin \varphi)$ ,  $\gamma_1(\varphi) = ab\mu_1(\varphi)D^{-2}\gamma_0(\varphi)$ , and  $D = \text{diag}(a, b)$  as above. The generating function of the billiard map inside the perturbed ellipse is

$$h_\epsilon(\varphi, \varphi') = |\gamma_\epsilon(\varphi') - \gamma_\epsilon(\varphi)| = h_0(\varphi, \varphi') + \epsilon h_1(\varphi, \varphi') + O(\epsilon^2).$$

The first terms of this expansion satisfy the identities  $h_0(\varphi, \varphi') = |\gamma_0(\varphi') - \gamma_0(\varphi)|$  and  $h_0(\varphi, \varphi')h_1(\varphi, \varphi') = \langle \gamma_0(\varphi') - \gamma_0(\varphi), \gamma_1(\varphi') - \gamma_1(\varphi) \rangle$ .

Let  $(q_j)_{j \in \mathbb{Z}}$  be the billiard trajectory inside the ellipse  $Q$  with caustic  $C_\lambda$  such that  $q_j = \gamma_0(\varphi_j)$  and  $\varphi_0 = \varphi$ . The unit inward velocities of this trajectory are

$$p_j = \frac{q_{j+1} - q_j}{|q_{j+1} - q_j|} = \frac{\gamma_0(\varphi_{j+1}) - \gamma_0(\varphi_j)}{h_0(\varphi_j, \varphi_{j+1})}.$$

It follows from Proposition 2.1 that the subharmonic Melnikov potential is

$$\begin{aligned} L_1(\varphi) &= \sum_{j=0}^{n-1} h_1(\varphi_j, \varphi_{j+1}) \\ &= \sum_{j=0}^{n-1} \langle p_j, \gamma_1(\varphi_{j+1}) - \gamma_1(\varphi_j) \rangle \\ &= ab \sum_{j=0}^{n-1} \langle p_j, \mu_1(\varphi_{j+1})D^{-2}q_{j+1} - \mu_1(\varphi_j)D^{-2}q_j \rangle \\ &= ab \sum_{j=0}^{n-1} \langle p_{j-1} - p_j, D^{-2}q_j \rangle \mu_1(\varphi_j) \\ &= 2\lambda \sum_{j=0}^{n-1} \mu_1(\varphi_j). \end{aligned}$$

We have used the periodicity and Lemma 4.1 in the last two equalities. □

Next, we give a couple of sufficient conditions for the subharmonic Melnikov potential to be constant. These conditions are trivial. Nevertheless, they play a key role in our problem. Concretely, we shall check later on that they are also necessary conditions in the class of  $2\pi$ -periodic entire functions  $\mu_1(\varphi)$ .

**COROLLARY 4.1.** *Let  $\mu_1(\varphi)$  be any  $2\pi$ -periodic smooth function.*

- (1) *If the period  $n$  is odd, then  $\mu_1(\varphi)$  constant  $\Rightarrow L_1(\varphi)$  constant.*
- (2) *If the period  $n$  is even, then  $\mu_1'(\varphi)$   $\pi$ -antiperiodic  $\Rightarrow L_1(\varphi)$  constant.*

*Proof.* The case  $n$  odd is obvious. If  $n$  is even, the  $(m, n)$ -gons inscribed in  $Q$  and circumscribed around  $C_\lambda$  are symmetric with respect to the origin, so  $\varphi_{j+n/2} = \varphi_j + \pi$  and

$$L'_1(\varphi) = 2\lambda \sum_{j=0}^{n-1} \mu'_1(\varphi_j) = 2\lambda \sum_{j=0}^{n/2-1} (\mu'_1(\varphi_j) + \mu'_1(\varphi_j + \pi)).$$

In particular,  $n$  even and  $\mu'_1(\varphi)$   $\pi$ -antiperiodic  $\Rightarrow L'_1(\varphi) \equiv 0 \Rightarrow L_1(\varphi)$  constant. □

The subharmonic Melnikov potential of the  $(m, n)$ -resonant caustic for the perturbed circle (2) is

$$L_1(\theta) = 2r_0 \sin(m\pi/n) \sum_{j=0}^{n-1} r_1(\theta_j), \quad \theta_j = \theta + 2\pi m j/n; \tag{8}$$

see [19, Proposition 10]. We recall that  $\lambda = r_0 \sin(m\pi/n)$  is the  $(m, n)$ -resonant caustic parameter of the circle of radius  $r_0$ . Furthermore, all the  $(m, n)$ -gons inscribed in the circle of radius  $r_0$  and circumscribed around the circle of radius  $\lambda = r_0 \sin(m\pi/n)$  are regular, so their vertices are of the form  $q_j = (r_0 \cos \theta_j, r_0 \sin \theta_j)$  with  $\theta_j = \theta + 2\pi m j/n$ . Hence, function (8) is the limit of function (7) when both  $a$  and  $b$  tend to  $r_0$ .

Although functions (7) and (8) look quite similar, they conceal a crucial difference. There is a simple formula for the  $\theta_j$  angles, but not for the  $\varphi_j$  ones. This has to do with the fact that the billiard trajectories inside a circle of radius  $r_0$  sharing a circular caustic with radius  $\lambda = r_0 \sin(\delta/2)$  have a rigid angular dynamics of the form  $\theta \mapsto \theta + \delta$ . In contrast, such a rigid angular dynamics does not take place for elliptic tables when the angle  $\varphi$  is considered, which is a source of technical difficulties in the study of the subharmonic Melnikov potential (7). Nevertheless, it is possible to define a new angular parameter  $t$  over the ellipse  $Q$  in such a way that all billiard trajectories inside  $Q$  sharing the elliptical caustic  $C_\lambda$  have a rigid angular dynamics of the form  $t \mapsto t + \delta$ , for some constant shift  $\delta = \delta(\lambda)$ .

We need some notation for elliptic functions in order to define this angular parameter  $t$ . We refer to [1, 22] for a general background on elliptic functions. Given a quantity  $k \in (0, 1)$ , called the *modulus*, the *complete elliptic integral of the first kind* is  $K = K(k) = \int_0^{\pi/2} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi$ . We also write  $K' = K'(k) = K(\sqrt{1 - k^2})$ . The *amplitude* function  $\varphi = \text{am } t$  is defined through the inversion of the integral

$$t = \int_0^\varphi (1 - k^2 \sin^2 \phi)^{-1/2} d\phi.$$

Then the *elliptic sine* and the *elliptic cosine* are defined by the trigonometric relations

$$\text{sn } t = \sin \varphi, \quad \text{cn } t = \cos \varphi,$$

respectively. Dependence on the modulus is denoted by a comma preceding it, so we can write  $\text{am}(t, k)$ ,  $\text{sn}(t, k)$ , and  $\text{cn}(t, k)$  to avoid any confusion. In the following lemma it is stated that the angular dynamics becomes rigid in the angular parameter  $t$  given by  $\varphi = \text{am}(t, k)$ . It suffices to find the suitable modulus  $k$  for each elliptical caustic  $C_\lambda$ .

LEMMA 4.2. *Having fixed a caustic parameter  $\lambda \in (0, b)$ , we set the modulus  $k \in (0, 1)$  and the constant shift  $\delta \in (0, 2K)$  by the formulae*

$$k^2 = \frac{a^2 - b^2}{a^2 - \lambda^2}, \quad \frac{\delta}{2} = \int_0^{\vartheta/2} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi, \tag{9}$$

where  $\vartheta \in (0, \pi)$  is the angle such that  $\sin(\vartheta/2) = \lambda/b$ . Let

$$q_j = (a \cos \varphi_j, b \sin \varphi_j) = (a \operatorname{cn}(t_j, k), b \operatorname{sn}(t_j, k))$$

be any billiard trajectory inside the ellipse  $Q$  with caustic  $C_\lambda$ . Then  $t_{j+1} = t_j + \delta$ .

*Proof.* By definition,  $\varphi_j = \operatorname{am}(t_j, k)$ , so  $t_{j+1} - t_j = \int_{\varphi_j}^{\varphi_{j+1}} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi$ . These integrals are equal to a constant  $\delta$  that depends only on  $C_\lambda$ ; see [5, p. 1543]. The formula for the constant shift is given in [5, p. 1540]. □

Observe that if  $a = b = r_0$  then the modulus  $k$  is equal to zero, the complete elliptic integral  $K$  is equal to  $\pi/2$ , the amplitude function is the identity, the elliptic sine/cosine are the usual sine/cosine, the shift  $\delta \in (0, \pi)$  is given by  $\lambda = r_0 \sin(\delta/2)$ , and the dynamical relation  $t_{j+1} = t_j + \delta$  becomes  $\varphi_{j+1} = \varphi_j + \delta$ . Thus, we recover the known rigid angular dynamics for circular tables as a limit of the formulae for elliptic tables.

From now on,  $k$  and  $\delta$  will denote the modulus and the constant shift defined in (9). Thus, we shall ignore the dependence of the elliptic functions on the modulus. We note that  $C_\lambda$  has eccentricity  $k$ . Furthermore,  $C_\lambda$  is the  $(m, n)$ -resonant elliptical caustic if and only if

$$n\delta = 4Km. \tag{10}$$

This identity has the following geometric interpretation. When a billiard trajectory makes one turn around  $C_\lambda$ , the old angular variable  $\varphi$  changes by  $2\pi$ , so the new angular variable  $t$  changes by  $4K$ . On the other hand, we have seen that the variable  $t$  changes by  $\delta$  when a billiard trajectory bounces once. Hence, a billiard trajectory inscribed in  $Q$  and circumscribed around  $C_\lambda$  makes exactly  $m$  turns around  $C_\lambda$  after  $n$  bounces if and only if (10) holds.

PROPOSITION 4.2. *Let  $\mu_1(\varphi)$  be any  $2\pi$ -periodic entire function.*

- (1) *If the period  $n$  is odd, then  $L_1(\varphi)$  constant  $\Leftrightarrow \mu_1(\varphi)$  constant.*
- (2) *If the period  $n$  is even, then  $L_1(\varphi)$  constant  $\Leftrightarrow \mu'_1(\varphi)$   $\pi$ -antiperiodic.*

*Proof.* Let  $\Delta = 2K + 2K'i$  and  $z(t) = \operatorname{cn} t + i \operatorname{sn} t$ . If  $\varphi = \operatorname{am} t$ , then

$$e^{i\varphi} = \cos \varphi + i \sin \varphi = \operatorname{cn} t + i \operatorname{sn} t = z(t),$$

$$e^{-i\varphi} = \cos \varphi - i \sin \varphi = \operatorname{cn} t - i \operatorname{sn} t = z(t + \Delta).$$

We have used the fact that the elliptic sine is  $\Delta$ -antiperiodic, and the elliptic cosine is  $\Delta$ -periodic. We also recall that the elliptic sine/cosine are  $2K$ -antiperiodic meromorphic functions on the whole complex plane whose unique singularities are the points of the form

$$\tau_{r,s} = 2Kr + (1 + 2s)K'i, \quad r, s \in \mathbb{Z}.$$

Furthermore, these singularities are just simple poles whose residues are

$$\operatorname{res}(\operatorname{cn}; \tau_{r,s}) = (-1)^{r+s+1}i/k, \quad \operatorname{res}(\operatorname{sn}; \tau_{r,s}) = (-1)^r/k.$$

Thus,  $z(t)$  is a  $2K$ -antiperiodic meromorphic function whose unique singularities are the points of the set

$$P = \{\tau_{r,2s+1} : r, s \in \mathbb{Z}\} = \tau_* + 2K\mathbb{Z} + 4K'i\mathbb{Z}, \quad \tau_* = \tau_{0,-1} = -K'i.$$

As before, these singularities are just simple poles.

Let  $\sum_{l \in \mathbb{Z}} \hat{\mu}_l e^{il\varphi}$  be the Fourier expansion of  $\mu_1(\varphi)$ . Then

$$\mu_1(\text{am } t) = \mu_1(\varphi) = \sum_{l \in \mathbb{Z}} \hat{\mu}_l e^{il\varphi} = \hat{\mu}_-(z(t + \Delta)) + \hat{\mu}_0 + \hat{\mu}_+(z(t)),$$

where  $\hat{\mu}_-(z) = \sum_{l=1}^{\infty} \hat{\mu}_{-l} z^l$  and  $\hat{\mu}_+(z) = \sum_{l=1}^{\infty} \hat{\mu}_l z^l$ . We note that the functions  $\hat{\mu}_{\pm}(z)$  are entire, because  $\mu_1(\varphi)$  is entire. Furthermore,

$$L_1(\text{am } t) = L_1(\varphi) = 2\lambda \sum_{j=0}^{n-1} \mu_1(\varphi_j) = 2\lambda(L_-(t) + n\hat{\mu}_0 + L_+(t)), \tag{11}$$

where

$$L_-(t) = \sum_{j=0}^{n-1} \hat{\mu}_-(z(t + \Delta + j\delta)) \quad \text{and} \quad L_+(t) = \sum_{j=0}^{n-1} \hat{\mu}_+(z(t + j\delta)).$$

Let us study the behaviour of these two functions around the point  $\tau_* = -K'i$ . Concretely, we shall prove that  $L_-(t)$  is analytic at  $t = \tau_*$ , whereas  $L_+(t)$  has a non-removable singularity at  $t = \tau_*$  provided  $\mu_1(\varphi)$  is non-constant and  $n$  is odd, or provided  $\mu'_1(\varphi)$  is not  $\pi$ -antiperiodic and  $n$  is even.

We begin with a couple of simple observations. If  $j \in \{0, \dots, n - 1\}$ , then:

- (a)  $\Im(\tau_* + \Delta + j\delta) = K'$ , so  $\tau_* + \Delta + j\delta \notin P$ ; and
- (b)  $\tau_* + j\delta \in P \Leftrightarrow 4Kmj/n = j\delta \in 2K\mathbb{Z} \Leftrightarrow 2jm \in n\mathbb{Z} \Leftrightarrow 2j \in n\mathbb{Z} \Leftrightarrow j \in \{0, n/2\}$ .  
Here, we have used the fact that  $\delta \in \mathbb{R}$ , equation (10), and  $\text{gcd}(m, n) = 1$ . Furthermore, we stress that the equality  $j = n/2$  can only occur when  $n$  is even.

We deduce the following results from the above observations.

- (1)  $L_-(t)$  is analytic at  $t = \tau_*$ , because so are  $z(t + \Delta + j\delta)$  for  $j = 0, \dots, n - 1$ .
- (2) If  $n$  is odd and  $\mu_1(\varphi)$  is non-constant, then:
  - the function  $\hat{\mu}_+(z)$  is non-constant and entire;
  - the function  $L_+(t) - \hat{\mu}_+(z(t)) = \sum_{j=1}^{n-1} \hat{\mu}_+(z(t + j\delta))$  is analytic at  $t = \tau_*$ ;
  - the composition  $\hat{\mu}_+(z(t))$  has a non-removable singularity at  $t = \tau_*$ ; and
  - the function (11) is non-constant, since it has a non-removable singularity at  $t = \tau_*$ .
- (3) If  $n$  is even and  $\mu'_1(\varphi)$  is not  $\pi$ -antiperiodic, then:
  - the sum  $\hat{\sigma}(z) = \hat{\mu}_+(z) + \hat{\mu}_+(-z) = 2 \sum_{l=1}^{\infty} \hat{\mu}_{2l} z^{2l}$  is a non-constant entire function;
  - $z(t + n\delta/2) = z(t + 2Km) = (-1)^m z(t) = -z(t)$ , since  $m$  is odd;
  - $\hat{\mu}_+(z(t)) + \hat{\mu}_+(z(t + n\delta/2)) = \hat{\sigma}(z(t))$ ;
  - the function  $L_+(t) - \hat{\sigma}(z(t))$  is analytic at  $t = \tau_*$ ;
  - the composition  $\hat{\sigma}(z(t))$  has a non-removable singularity at  $t = \tau_*$ ; and
  - the function (11) is non-constant, since it has a non-removable singularity at  $t = \tau_*$ .

Therefore, the proof follows by combining the above results with Corollary 4.1. □

Finally, we note that our main result (Theorem 1.1) follows directly from Corollary 3.1 and Proposition 4.2.

*Acknowledgements.* The first author was partially supported by the CRM (Catalonia) and Brazilian agencies CNPq and FAPEMIG. The second author was supported in part by MICINN-FEDER grant MTM2009-06973 (Spain) and CUR-DIUE grant 2009SGR859 (Catalonia). This work was completed while the first author was a visitor at the CRM at Barcelona (Spain). Useful conversations with Pablo S. Casas, Amadeu Delshams, Vadim Kaloshin, and Vassilios Rothos are gratefully acknowledged.

## REFERENCES

- [1] M. Abramowitz and I. Stegun. *Handbook of Mathematical Functions*. Dover, New York, 1972.
- [2] Yu. Baryshnikov and V. Zharnitsky. Sub-Riemannian geometry and periodic orbits in classical billiards. *Math. Res. Lett.* **13** (2006), 587–598.
- [3] G. D. Birkhoff. *Dynamical Systems (American Mathematical Society Colloquium Publications)*. American Mathematical Society, Providence, RI, 1966, original ed. 1927.
- [4] P. S. Casas and R. Ramírez-Ros. The frequency map for elliptic billiards. *SIAM J. Appl. Dyn. Syst.* **10** (2011), 278–324.
- [5] S.-J. Chang and R. Friedberg. Elliptical billiards and Poncelet’s theorem. *J. Math. Phys.* **29** (1988), 1537–1550.
- [6] A. Delshams and R. Ramírez-Ros. Poincaré–Melnikov–Arnold method for analytic planar maps. *Nonlinearity* **9** (1996), 1–26.
- [7] R. Douady. *Applications du théorème des tores invariants*. Thèse de 3ème Cycle, Univ. Paris VII, 1982.
- [8] V. Dragović and M. Radnović. Bifurcations of Liouville tori in elliptical billiards. *Regul. Chaotic Dyn.* **14** (2009), 479–494.
- [9] V. Dragović and M. Radnović. *Poncelet Porisms and Beyond*. Birkhäuser, Basel, 2011.
- [10] E. Gutkin and A. Katok. Caustics for inner and outer billiards. *Comm. Math. Phys.* **173** (1995), 101–133.
- [11] E. Gutkin. Billiard dynamics: a survey with the emphasis on open problems. *Regul. Chaotic Dyn.* **8** (2003), 1–13.
- [12] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, Cambridge, 1995.
- [13] O. Knill. On nonconvex caustics of convex billiards. *Elem. Math.* **53** (1998), 89–106.
- [14] V. V. Kozlov and D. Treshchëv. *Billiards: A Genetic Introduction to the Dynamics of Systems with Impacts (Translations of Mathematical Monographs, 89)*. American Mathematical Society, Providence, RI, 1991.
- [15] V. F. Lazutkin. The existence of caustics for a billiard problem in a convex domain. *Math. USSR Izvestija* **7** (1973), 185–214.
- [16] J. Mather. Glancing billiards. *Ergod. Th. & Dynam. Sys.* **2** (1982), 397–403.
- [17] J. D. Meiss. Symplectic maps, variational principles, and transport. *Rev. Mod. Phys.* **64** (1992), 795–848.
- [18] J. V. Poncelet. *Traité des Propriétés Projectives des Figures*. Gauthier-Villars, Paris, 1866.
- [19] R. Ramírez-Ros. Break-up of resonant invariant curves in billiards and dual billiards associated to perturbed circular tables. *Physica D* **214** (2006), 78–87.
- [20] V. Rothos. Subharmonic bifurcations of localized solutions of a discrete NLS equation. *Discrete Contin. Dyn. Syst.* (2005), 756–767, supplement volume.
- [21] S. Tabachnikov. *Billiards (Panorama et Synthèses)*. Société Mathématique de France, Paris, 1995.
- [22] E. T. Whittaker and G. N. Watson. *A Course of Modern Analysis*. Cambridge University Press, Cambridge, 1927.